REPORT DOCUMENTATION PAGE

Form Approved OMB NO. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggesstions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any oenalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	DATE (DD-MM-YYYY) 2. REPORT TYPE		3. DATES COVERED (From - To)		
	New Reprint		-		
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER			
Genome Sequences of Four Acinetobacter baumannii-A.			W911NF-11-1-0136		
calcoaceticus Complex Isolates from Combat-Related Infections Sustained in the Middle East		5b. GRANT NUMBER			
			5c. PROGRAM ELEMENT NUMBER		
			206022		
6. AUTHORS			5d. PROJECT NUMBER		
P. Ketter, M. N. Guentzel, J. P. Chambers, J. Jorgensen, C. K. Murray,					
A. P. Cap, JJ. Yu, M. Eppinger, B. P. Arulanandam		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES			8. PERFORMING ORGANIZATION REPORT NUMBER		
University of Texas at San Antonio One UTSA Circle					
San Antonio, TX 782	49 -0603				
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS (ES)			10. SPONSOR/MONITOR'S ACRONYM(S) ARO		
U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
			59027-LS-REP.64		
12. DISTRIBUTION AVAILIBILITY STAT	EMENT				

Approved for public release; distribution is unlimited.

13. SUPPLEMENTARY NOTES

The views, opinions and/or findings contained in this report are those of the author(s) and should not contrued as an official Department of the Army position, policy or decision, unless so designated by other documentation.

14. ABSTRACT

Acinetobacter baumannii is among the most prevalent bacterial causes of combat-related infections on the battlefield. Antibiotic resistance and a poor understanding of the protective host immune responses make treatment difficult. Here, we report the genome sequences of four clinical Acinetobacter baumannii- A. calcoaceticus complex isolates exhibiting significant differences in virulance in a mouse sepsis model.

15. SUBJECT TERMS

Acinetobacter baumannii, A. calcoaceticus, combat-related infections

16. SECURITY CLASSIFICATION OF:		17. Emiliarion or	I C. T.OTHEBETT	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE	ABSTRACT	OF PAGES	Bernard Arulanandam
UU	UU	υυ	UU		19b. TELEPHONE NUMBER 210-458-5492

Report Title

Genome Sequences of Four Acinetobacter baumannii-A. calcoaceticus Complex Isolates from Combat-Related Infections Sustained in the Middle East

ABSTRACT

Acinetobacter baumannii is among the most prevalent bacterial causes of combat-related infections on the battlefield. Antibiotic resistance and a poor understanding of the protective host immune responses make treatment difficult. Here, we report the genome sequences of four clinical Acinetobacter baumannii- A. calcoaceticus complex isolates exhibiting significant differences in virulance in a mouse sepsis model.

REPORT DOCUMENTATION PAGE (SF298) (Continuation Sheet)

Continuation for Block 13

ARO Report Number 59027.64-LS-REP Genome Sequences of Four Acinetobacter baur...

Block 13: Supplementary Note

© 2014 . Published in Genome Announcements, Vol. Ed. 0 2, (1) (2014), (, (1). DoD Components reserve a royalty-free, nonexclusive and irrevocable right to reproduce, publish, or otherwise use the work for Federal purposes, and to authroize others to do so (DODGARS §32.36). The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.

Approved for public release; distribution is unlimited.



Genome Sequences of Four Acinetobacter baumannii-A. calcoaceticus Complex Isolates from Combat-Related Infections Sustained in the Middle East

Patrick Ketter, M. Neal Guentzel, James P. Chambers, James Jorgensen, Clinton K. Murray, Andrew P. Cap, Jieh-Juen Yu, Andrew P. Cap, Jieh-Juen Yu, Andrew P. Cap, Lands Jorgensen, Clinton K. Murray, Lands Jorgensen, Lands Jorgen Mark Eppinger, Bernard P. Arulanandama

University of Texas at San Antonio, San Antonio, Texas, USA^a; University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA^b; San Antonio Military Medical Center, JBSA-Fort Sam Houston, San Antonio, Texas, USA-Fort Sam Houston, San Antonio Military Medical Center, JBSA-Fort Sam Houston San Antonio Texas USAd

Acinetobacter baumannii is among the most prevalent bacterial causes of combat-related infections on the battlefield. Antibiotic resistance and a poor understanding of the protective host immune responses make treatment difficult. Here, we report the genome sequences of four clinical Acinetobacter baumannii-A. calcoaceticus complex isolates exhibiting significant differences in virulence in a mouse sepsis model.

Received 10 January 2014 Accepted 15 January 2014 Published 6 February 2014

Citation Ketter P, Guentzel MN, Chambers JP, Jorgensen J, Murray CK, Cap AP, Yu J-J, Eppinger M, Arulanandam BP. 2014. Genome sequences of four Acinetobacter baumannii-A. calcoaceticus complex isolates from combat-related infections sustained in the Middle East. Genome Announc. 2(1):e00026-14. doi:10.1128/genomeA.00026-14.

Copyright © 2014 Ketter et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Address correspondence to Bernard P. Arulanandam, bernard.arulanandam@utsa.edu.

cinetobacter baumannii accounts for >36% of combat-related infections resulting from injuries sustained by military service personnel in the Middle East, making it among the most prevalent bacterial pathogens encountered on the battlefield (1, 2). Ubiquitous in nature, A. baumannii resists desiccation, allowing the pathogen to survive in relatively arid environments most other bacteria cannot (3-5). This, in combination with the increased level of antibiotic resistance associated with A. baumannii and its ability to colonize various bodily sites, makes these infections difficult to control and treat (3,6–11). The limited genomic plasticity and biochemical conservation among Acinetobacter species pose problems with regard to species identification in clinical settings, resulting in A. baumannii and Acinetobacter calcoaceticus often being reported as the A. baumannii-A. calcoaceticus complex (ABC) (10, 12). Genomic heterogeneity in A. baumannii has an established impact on patient outcome based on variations of the virulence and antibiotic resistance gene alleles present in each strain (13-16).

To this end, we sequenced four clinical multidrug-resistant ABC strains, CI77, CI78, CI79, and CI86, from a collection obtained from respiratory (CI78 and CI79) or wound (CI77 and CI86) cultures from military personnel injured in either Iraq or Afghanistan (17). In a murine sepsis model, we observed significant differences in virulence in vivo between strains CI77 and CI79 correlating to sustained and elevated blood serum levels of an acute-phase small pattern recognition receptor known as PTX3; PTX3 possesses antimicrobial properties following challenge with strain CI79 (P. M. Ketter, M. N. Guentzel, B. Schaffer, M. Herzig, X. Wu, C. G. Fedyk, J. Yu, J. Jorgensen, J. P. Chambers, A. P. Cap, and B. P. Arulanandam, submitted for publication). Genomic DNA was subjected to Illumina sequencing using paired-end libraries with 300-bp inserts on the HiSeq 2000 platform. The draft genome was assembled with the Velvet assembler (18, 19), and the IGS Annotation Engine and Manatee were used for structural and functional genome annotation and visualization of the chromosomes and plasmid contigs (20).

The observed genome sizes of the isolates range from 3.8 to 4.2 Mb, with an average G+C content of 38%. In silico comparison of the 16s rRNA genes allowed the complex and species attribution of C177, C179, and C186 as being pathogenic A. baumannii strains, while C178 is representative of A. calcoaceticus, is generally considered nonpathogenic, and is found to occasionally colonize but rarely cause infection in humans (10, 12). We determined the carriage of several reported A. baumannii virulence factors, such as LpxC, PglL, and AbaI (21-23), and found only limited genomic plasticity among the analyzed Middle Eastern ABC isolates. The availability of these sequences as a reference in RNAseq analysis will assist in future studies examining alterations in the ABC complex transcriptome and will help enhance our understanding of the underlying genetics and regulatory pathways leading to altered clinical disease manifestation in patients infected with emerging pathogens of ABC.

Nucleotide sequence accession numbers. The genome sequences of the Acinetobacter clinical isolates C177, C178, C179, and C186 have been deposited in GenBank under accession no. AVOC00000000, AVOE00000000, AVOD00000000, and AVOB00000000, respectively.

ACKNOWLEDGMENTS

This research was partially funded by the Army Research Office of the Department of Defense (contract no. W911NF-11-1-0136) and the Álvarez Graduate Research Education Excellence Fund. Computational support was provided by the Computational System Biology Core at the University of Texas at San Antonio and the South Texas Center for Emerging Infectious Diseases (STCEID).

The opinions or assertions contained herein are the private views of

the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

REFERENCES

- Petersen K, Riddle MS, Danko JR, Blazes DL, Hayden R, Tasker SA, Dunne JR. 2007. Trauma-related infections in battlefield casualties from Iraq. Ann. Surg. 245:803–811. http://dx.doi.org/10.1097/01.sla.00002517 07.32332.cl.
- Tribble DR, Conger NG, Fraser S, Gleeson TD, Wilkins K, Antonille T, Weintrob A, Ganesan A, Gaskins LJ, Li P, Grandits G, Landrum ML, Hospenthal DR, Millar EV, Blackbourne LH, Dunne JR, Craft D, Mende K, Wortmann GW, Herlihy R, McDonald J, Murray CK. 2011. Infection-associated clinical outcomes in hospitalized medical evacuees after traumatic injury: trauma infectious disease outcome study. J. Trauma 71:S33—S42. http://dx.doi.org/10.1097/TA.0b013e318221162e.
- Bergogne-Bérézin F, Towner KJ. 1996. Acinetobacter spp. as nosocomial pathogens: microbiological, clinical, and epidemiological features. Clin. Microbiol. Rev. 9:148–165.
- Fournier PE, Richet H. 2006. The epidemiology and control of *Acineto-bacter baumannii* in health care facilities. Clin. Infect. Dis. 42:692–699. http://dx.doi.org/10.1086/500202.
- Baumann P. 1968. Isolation of Acinetobacter from soil and water. J. Bacteriol. 96:39

 –42.
- Koeleman JG, van der Bijl MW, Stoof J, Vandenbroucke-Grauls CM, Savelkoul PH. 2001. Antibiotic resistance is a major risk factor for epidemic behavior of Acinetobacter baumannii. Infect. Control Hosp. Epidemiol. 22:284–288. http://dx.doi.org/10.1086/501901.
- Cisneros JM, Reyes MJ, Pachón J, Becerril B, Caballero FJ, García-Garmendia JL, Ortiz C, Cobacho AR. 1996. Bacteremia due to Acinetobacter baumannii: epidemiology, clinical findings, and prognostic features. Clin. Infect. Dis. 22:1026–1032. http://dx.doi.org/10.1093/clinids/22.6.1026.
- Dy ME, Nord JA, LaBombardi VJ, Kislak JW. 1999. The emergence of resistant strains of *Acinetobacter baumannii*: clinical and infection control implications. Infect. Control Hosp. Epidemiol. 20:565–567. http://dx.doi .org/10.1086/501673.
- Abbo A, Navon-Venezia S, Hammer-Muntz O, Krichali T, Siegman-Igra Y, Carmeli Y. 2005. Multidrug-resistant Acinetobacter baumannii. Emerg. Infect. Dis. 11:22–29. http://dx.doi.org/10.3201/eid1101.040001.
- Peleg AY, de Breij A, Adams MD, Cerqueira GM, Mocali S, Galardini M, Nibbering PH, Earl AM, Ward DV, Paterson DL, Seifert H, Dijkshoorn L. 2012. The success of *Acinetobacter* species; genetic, metabolic and virulence attributes. PLoS One 7:e46984. http://dx.doi.org/10.1371 /journal.pone.0046984.
- Hospenthal DR, Crouch HK, English JF, Leach F, Pool J, Conger NG, Whitman TJ, Wortmann GW, Robertson JL, Murray CK. 2011. Multidrug-resistant bacterial colonization of combat-injured personnel at admission to medical centers after evacuation from Afghanistan and Iraq. J. Trauma 71:S52–S57. http://dx.doi.org/10.1097/TA.0b013e31822118fb.
- Chiang MC, Kuo SC, Chen YC, Lee YT, Chen TL, Fung CP. 2011.
 Polymerase chain reaction assay for the detection of Acinetobacter bau-

- mannii in endotracheal aspirates from patients in the intensive care unit. J. Microbiol. Immunol. Infect. 44:106–110. http://dx.doi.org/10.1016/j.jmii .2010.04.003.
- Sahl JW, Johnson JK, Harris AD, Phillippy AM, Hsiao WW, Thom KA, Rasko DA. 2011. Genomic comparison of multi-drug resistant invasive and colonizing *Acinetobacter baumannii* isolated from diverse human body sites reveals genomic plasticity. BMC Genomics 12:291. http://dx .doi.org/10.1186/1471-2164-12-291.
- Kang G, Hartzell JD, Howard R, Wood-Morris RN, Johnson MD, Fraser S, Weintrob A, Wortmann G. 2010. Mortality associated with Acinetobacter baumannii complex bacteremia among patients with warrelated trauma. Infect. Control Hosp. Epidemiol. 31:92–94. http://dx.doi .org/10.1086/649220.
- Johnson EN, Burns TC, Hayda RA, Hospenthal DR, Murray CK. 2007. Infectious complications of open type III tibial fractures among combat casualties. Clin. Infect. Dis. 45:409–415. http://dx.doi.org/10 .1086/520029.
- Albrecht MC, Albrecht MA, Griffith ME, Murray CK, Chung KK, Horvath EE, Ward JA, Hospenthal DR, Holcomb JB. 2006. Impact of Acinetobacter infection on the mortality of burn patients. J. Am. Coll. Surg. 203:546–550. http://dx.doi.org/10.1016/j.jamcollsurg.2006.06.013.
- Hawley JS, Murray CK, Griffith ME, McElmeel ML, Fulcher LC, Hospenthal DR, Jorgensen JH. 2007. Susceptibility of *Acinetobacter* strains isolated from deployed U.S. military personnel. Antimicrob. Agents Chemother. 51:376–378. http://dx.doi.org/10.1128/AAC.00858-06.
- Zerbino DR. 2002. Using the Velvet de novo assembler for short-read sequencing technologies, Curr. Protoc. Bioinformatics 31:11.5.1–11.5.12. http://dx.doi.org/10.1002/0471250953.bi1105s31.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. Genome Res. 18:821–829. http://dx.doi .org/10.1101/gr.074492.107.
- Galens K, Orvis J, Daugherty S, Creasy HH, Angiuoli S, White O, Wortman J, Mahurkar A, Gwinn Giglio M. 2011. The IGS standard operating procedure for automated prokaryotic annotation. Stand. Genomic Sci. 4:244–251. http://dx.doi.org/10.4056/sigs.1223234.
- Iwashkiw JA, Seper A, Weber BS, Scott NE, Vinogradov E, Stratilo C, Reiz B, Cordwell SJ, Whittal R, Schild S, Feldman MF. 2012. Identification of a general O-linked protein glycosylation system in Acinetobacter baumannii and its role in virulence and biofilm formation. PLoS Pathog. 8:e1002758. http://dx.doi.org/10.1371/journal.ppat.1002758.
- Lin L, Tan B, Pantapalangkoor P, Ho T, Baquir B, Tomaras A, Montgomery JI, Reilly U, Barbacci EG, Hujer K, Bonomo RA, Fernandez L, Hancock RE, Adams MD, French SW, Buslon VS, Spellberg B. 2012. Inhibition of LpxC protects mice from resistant *Acinetobacter baumannii* by modulating inflammation and enhancing phagocytosis. mBio 3(5): e00312-12. http://dx.doi.org/10.1128/mBio.00312-12.
- Niu C, Clemmer KM, Bonomo RA, Rather PN. 2008. Isolation and characterization of an autoinducer synthase from *Acinetobacter bauman*nii. J. Bacteriol. 190:3386–3392. http://dx.doi.org/10.1128/JB.01929-07.